

Nuevo tratamiento de reducción de volumen endoscópico en enfisema agudo

Zephyr®



CRITERIOS DE INCLUSION

- ‡ 'LVQHD \ R OLPLWDFLyQ D ODV WDUHDV KDE WUDWDPLHQWR ySWLPR
- ‡ 'LDJQRVWLFRR FOfQLFR GH HQILVHPD KHWHU
- ‡ &HVDFLyQ WDEiTXLFD GH DO PHQRV PHVHV
- ‡ 9ROXPHQ UHVLGXDO !
- ‡)(9
- ‡ '/&2F HQ SDFLHQWHV FRQ HQILVHPD KHV
- ‡ /D KLSHULQVXIOfLYQ GLQiPLFD SXHGH VHU SDUD OD WRPD GH GHFLVLYQ

CRITERIOS DE EXCLUSION

- ‡)XPDGRU DFWXDO
- ‡ 6HYHUD KLSHUVHFUHFLyQ GH PRFR
- ‡ %URQFR HVSDVPR PDUFDGR
- ‡ &RPRUELGDGHV DVRFLDGDV FRQ SURQyVWLI
- ‡ (QILVHPD KRPRJpQHR
- ‡ &RPRUELGDGHV TXH FDXVHQ GLVQHD LQVX KLSHUWHQVLYQ DQHPLD HWF

ESTUDIOS INICIALES REQUERIDOS

- ‡ ,PiJHQHV WRUiFLFDV
- ‡ 5D\RV [7yUD[7\$& 7\$& FRQ SURWRFROR 9,'
- ‡ 7HVW GH IXQFLyQ SXOPRQDU
- ‡ 7HVW GH PDUFKD GH PLQXWRV
- ‡ (FR FDUGLRJUDPD GRSSOHU
- ‡ *DVHV DUWHULDOHV
- ‡ 'RVDMH GH DOID DQWL WULSVLQD
- ‡ &XHVWLRQDULR GH 4/ 6W *HRUJH

pulmonX
Interventional Pulmonology

